

---

# Synthesis, Evaluation of Structure and Antimicrobial Activity of Metal Complexes of Schiff Bases Derived from Sulphonamide

**Anshika Hajela**

Deptt. of Chemistry, BMS College for Women, Bengaluru

**K. Chaturvedi**

Deptt. of Chemistry, Agra College, Agra

## ABSTRACT

Some new Cu (II), Zn (II), Mn (II) and Cr (III) complexes of Schiff bases derived from sulphonamide have been synthesized. They have been characterized on the basis of elemental analysis, IR and <sup>1</sup>HNMR spectroscopy. The spectral data suggests that Schiff base acts as bidentate ligand and an octahedral environment exist around the metal ion. In search for better fungicides and bactericides, biocidal studies were conducted against the organisms *Escherichia coli*, *Salmonella typhi*, *Bacillus subtilis*, *Staphylococcus aureus*, *Aspergillus flavous*, *Aspergillus niger*, *Penicillium citrinum* and *Fusarium oxysporum*. The result suggests that the metal complexes of Schiff base proved to be excellent bactericides and fungicides in the present investigations.

**KEYWORDS:** Cu (II), Zn (II), Mn (II) and Cr (III) complexes, IR and <sup>1</sup>HNMR spectroscopy and biocidal studies.

## INTRODUCTION

Sulphonamide are a group of compound which show distinct physical, chemical and biological properties.<sup>1-3</sup> Many chemotherapeutically important sulpha drugs like sulphadiazine, sulphathiazole, sulphamerazine and sulphonamide possess SO<sub>2</sub>NH<sub>2</sub> moiety which has an important toxicophoric function.<sup>4-6</sup> The Schiff bases synthesised from the sulpha drug possess pronounced antimicrobial<sup>7</sup>, antitumour<sup>8-9</sup>, antifouling<sup>10</sup>, antioncogenesis<sup>11</sup> activities. The condensation products of sulpha drugs with aldehydes and ketones are biologically active and also have good complexing ability; their activity increases on complexation with metal ions<sup>12-15</sup>.

The pronounced activity of metal complexes of Schiff bases derived from sulpha drugs has led to considerable interest in their coordination chemistry. A survey of literature reveals that very little attention has been paid on coordination behaviour of Schiff bases derived from sulpha drugs. The newly prepared complexes of sulpha drugs were characterized on the basis of elemental analysis and spectral studies. These complexes were also screened for their fungicidal and bactericidal activities.

## EXPERIMENTAL

All the chemicals used were of AR grade. The liquid reagents were purified by distillation.

p'-Benzylidene-p-aminobenzene sulphonamide (SA) was synthesized<sup>16</sup> by refluxing on water bath Sulphonamide (17.2g, 0.1 mole) and Benzaldehyde (10.6g, 0.1 mole) in 75 ml alcohol for two hours. On cooling the Schiff base SA separated out as white shining crystals. It was filtered, washed, dried and finally recrystallised from hot ethanol solution.

The Cu (II), Zn (II), Mn (II) and Cr (III) complexes of Schiff base were prepared<sup>17</sup> by refluxing on water bath, a mixture of 0.01 mole of metal chloride (i.e. 1.70 g of CuCl<sub>2</sub>. 2H<sub>2</sub>O, 1.36 g of ZnCl<sub>2</sub>, 1.97 g of MnCl<sub>2</sub>. 4H<sub>2</sub>O and 2.66 g of CrCl<sub>3</sub>. 6H<sub>2</sub>O) in 15 ml ethanol respectively with 0.02 mole of Schiff base SA (i.e. 5.2 g of SA)

in 20 ml ethanol for 3 hours. The reaction mixture was concentrated and then cooled. The solid derivatives were separated out. These were filtered, washed and finally air-dried.

For the microanalysis of C, H and N, CHN Perkin-Elmer micro analyzer 240 was used. The metal contents of the complexes were analyzed by standard methods. The IR spectra of the Schiff base and its metal complexes were recorded on Perkin-Elmer 4250 spectrophotometer in the range 4000-200  $\text{cm}^{-1}$  in CSI/KBr matrix. The  $^1\text{H}$ NMR spectra of the Schiff base and the metal complexes were recorded in  $\text{CDCl}_3$  on a Bruker DRX 300F, 300 MHz FTNMR spectrometer.

For the microanalysis of C, H and N, CHN Perkin-Elmer micro analyzer 240 was used. The metal contents of the complexes were analyzed by standard methods. The IR spectra of the Schiff base and its metal complexes were recorded on Perkin-Elmer 4250 spectrophotometer in the range 4000-200  $\text{cm}^{-1}$  in CSI/KBr matrix. The  $^1\text{H}$ NMR spectra of the Schiff base and the metal complexes were recorded in  $\text{CDCl}_3$  on a Bruker DRX 300F, 300 MHz FTNMR spectrometer.

### Antibacterial Activity

The bactericidal activity was evaluated by the filter paper disc diffusion method<sup>18</sup>. The nutrient agar medium (peptone, beef extract, NaCl and agar-agar) and 5mm diameter paper disc of Whatman No.1 were used. The compounds were dissolved in DMF in 100, 250 and 500 ppm concentrations. The filter paper discs were soaked in different solutions of the petridishes already seeded with the test organisms. The plates were incubated for 24 hours at 37°C and the inhibition zone around each disc was measured.

### Antifungal Activity

The fungicidal activity was evaluated by Agar plate technique<sup>19</sup>. The fungi were grown in agar medium (glucose 20g, starch 20g, agar-agar 20g and 1000ml water) at  $28 \pm 2^\circ\text{C}$  and the compounds after being dissolved in DMF were mixed in the medium. The growth of the fungus was obtained by measuring the diameter of colony in petridishes after 3 days and the percentage inhibition was calculated by the formula.

$$\% \text{ inhibition} = \frac{(C - T) \times 100}{C}$$

Where C and T are the diameters of the fungus colony in control and test plate, respectively.

## RESULTS AND DISCUSSION

The obtained metal complexes of the Schiff base are coloured solid, insoluble in water but soluble in benzene, acetone, DMF and other polar organic solvents. Elemental analyses suggests 1:2 (M:L) Stoichiometry<sup>20</sup>. All the physical and analytical results are listed in Table-1.

**Table-1. Physical & Analytical Data of Schiff base (SA) and its metal complexes**

COMPOUND	MOL.WT.	COLOUR	M.P. (°C)	ELEMENTAL ANALYSIS %				
				FOUND (CALCULATED)				
				C	H	N	S	METAL
SA [C <sub>13</sub> H <sub>12</sub> O <sub>2</sub> N <sub>2</sub> S]	260	White	162°-164°	60.10 (60.00)	4.50 (4.61)	10.72 (10.77)	12.25 (12.30)	- -
Cu-SA [Cu(C <sub>26</sub> H <sub>24</sub> N <sub>4</sub> O <sub>4</sub> S <sub>2</sub> ). 2H <sub>2</sub> O]	619	Brown	140°C	50.22 (50.40)	4.61 (4.52)	9.10 (9.04)	10.14 (10.33)	10.20 (10.17)
Zn-SA [Zn (C <sub>26</sub> H <sub>24</sub> N <sub>4</sub> O <sub>4</sub> S) 2H <sub>2</sub> O]	621	Cream	157°C	50.45 (50.24)	4.33(4 .50)	9.17 (9.01)	10.41 (10.30)	10.32 (10.46)
Mn-SA [Mn(C <sub>26</sub> H <sub>24</sub> N <sub>4</sub> O <sub>4</sub> S). 2H <sub>2</sub> O]	611	Light Brown	165°C	51.17 (51.06)	4.34 (4.58)	9.11 (9.16)	10.24 (10.47)	9.28 (9.00)
Cr-SA [Cr(C <sub>26</sub> H <sub>24</sub> N <sub>4</sub> O <sub>4</sub> S). 2H <sub>2</sub> O]	608	Light Green	173°C	51.45 (51.31)	4.57 (4.60)	9.41 (9.21)	10.34 (10.52)	8.67 (8.55)

### Infra Red Spectra

The Infra red spectra<sup>21</sup> of the Schiff base and the metal complexes show sharp bands in the region 3250-3300  $\text{cm}^{-1}$  and 1570-1590  $\text{cm}^{-1}$  attributed to the stretching and bending vibrations respectively of  $(\text{NH}_2)$ . While the symmetric and asymmetric vibrations of  $(\text{SO}_2)$  are observed in the region 1150-1160  $\text{cm}^{-1}$  and 1310-1330  $\text{cm}^{-1}$  respectively. The IR spectra of Schiff base show a sharp band at 1621  $\text{cm}^{-1}$  attributed to  $(\text{C}=\text{N})$  azomethine linkage<sup>21-23</sup>, a shift of  $\pm 15 \text{ cm}^{-1}$  is observed in this frequency in case of metal complexes which is suggestive of the coordination of the metal ion with the azomethine linkage. The band in the far IR spectra of metal complexes appear in the region 540-560  $\text{cm}^{-1}$  are tentatively assigned to  $(\text{M}-\text{N})$  vibrations. The stretching and bending vibrations found in the region 660-690  $\text{cm}^{-1}$  and 830-860  $\text{cm}^{-1}$  respectively attributed to the coordination of water molecules. The IR-spectral data is tabulated in Table-2.

**Table-2. Tentative assignments of some selected bands in the IR spectra ( $\text{cm}^{-1}$ ) of Schiff base (SA) and its metal complexes**

Compounds	$(\text{NH}_2)$ (stretch.)	$(\text{NH}_2)$ (bend.)	$(\text{SO}_2)$ (sym.)	$(\text{SO}_2)$ (asym)	$(\text{C}=\text{N})$	$(\text{H}_2\text{O})$ (stretch)	$(\text{H}_2\text{O})$ (bend.)	$(\text{M}-\text{N})$
SA	3295	1579	1154	1333	1621	-	-	-
Cu-SA	3251	1598	1169	1318	1645	679	833	550
Zn-SA	3296	1578	1155	1335	1628	622	840	555
Mn-SA	3293	1573	1154	1334	1631	683	838	551
Cr-SA	3303	1593	1161	1336	1627	627	834	549

### <sup>1</sup>HNMR Spectra

The <sup>1</sup>HNMR spectra of the Schiff base and its metal complexes show multiplet signals in the region 6.65-8.40 ppm due to benzene ring protons. While the signals due to  $\text{HC}=\text{N}$  protons<sup>23</sup> appear at 4.76 ppm in the spectra of Schiff base while it shifts downfield in the spectra of corresponding metal complexes. The broad signals due to  $-\text{NH}_2$  protons also shift downfield in the spectra of metal complexes. Thus, this downfield shift is attributed to the complex formation i.e. attachment of metal with nitrogen atom of  $-\text{NH}_2$  group and nitrogen atom<sup>24, 25</sup> of  $\text{HC}=\text{N}$  group. The evidences mentioned above supports the complex formation of the Schiff base. The <sup>1</sup>HNMR data is tabulated in Table-3.

**Table-3. <sup>1</sup>HNMR Spectral data of Schiff base (SA) and its metal complexes**

S.No.	Compound	Chemical Shift (ppm)	Peak position	Group Assigned.
1.	SA	(a) 6.65-8.40 (b) 4.89 (c) 10.02	Multiplet Singlet Singlet	Aromatic ring $\text{HC}=\text{N}$ $-\text{NH}_2$
2.	Cu-SA	(a) 6.54 -8.22 (b) 4.96 (c) 10.13	Multiplet Singlet Singlet	Aromatic ring $\text{HC}=\text{N}$ $-\text{NH}_2$
3.	Zn-SA	(a) 6.57-7.45 (b) 4.94 (c) 10.14	Multiplet Singlet Singlet	Aromatic ring $\text{HC}=\text{N}$ $-\text{NH}_2$
4.	Mn-SA	(a) 6.46-8.31 (b) 4.98 (c) 10.06	Multiplet Singlet Singlet	Aromatic ring $\text{HC}=\text{N}$ $-\text{NH}_2$
5.	Cr-SA	(a) 6.36-8.34 (b) 4.92 (c) 10.11	Multiplet Singlet Singlet	Aromatic ring $\text{HC}=\text{N}$ $-\text{NH}_2$

On the basis of spectroscopic characterization the tentative structure of Schiff base (SA) and its metal complexes is given in Figure-1 and Figure-2.




Fig 1 : p'- Benzylidene-p- aminobenzene sulphonamide (SA)




Fig 2: Tentative structure of the metal complexes


where  $M^n = \text{Cu (II), Zn (II), Mn (II) and Cr (III)}$  metal ions

LIGAND = Schiff Base SA

### Antibacterial and Antifungal Activities

The Schiff base (SA) as well as the metal complexes exhibited good amount of activity against all the tested bacteria and fungi. The Schiff base (SA) along with its metal complexes showed maximum activity against the bacteria, *E. coli* and minimum against the bacteria, *S. aureus*. Though, the fungicidal studies reveal that the activity was maximum against *A. niger* followed by *F. oxysporum*, *A. flavous* and *P. citrinum*.

The metal complexes were found to be more toxic as compared to their parent Schiff base. In general the activity of the metal complexes was in following order:-



The increased activity of metal complexes may be due to the effect of the metal ion configuration and the charge on normal cell. A possible mode of toxicity may be specified by chelation Theory. Chelation considerably reduces the polarity of the metal ion mainly because of partial sharing of its  $\pi$ -electrons and delocalization over the whole chelate ring. Such chelation increases the lipophilic character of metal chelate, which probably tends to break down the permeability barriers of cells, resulting in the interference with the normal cell process. Thus, the result suggests that the metal complexes of Schiff base (SA) have proved to be excellent bactericides and fungicides in the present investigations.

### REFERENCES

1. S Shah, R Vyas and RH Mehta. 1992, Ind. Chem. Soc.
2. HC Rani. 2005.J. Indian Council Chem.
3. SN Pandeya, D Sriram and G Nath. 1999. Eur. J. Pharma, Soc.
4. Wilson and Gisnold's R.F. Deorge (Ed.), J.B. Lippincott., Philadelphia, 1982, *Textbook of Organic Medicine and Pharmaceutical Chemistry*, 8<sup>th</sup> Edn.
5. G Kanagraj and GN Rao, 1993. Indian J. Chem.
6. SA Tilles, 2001 Southern Medical Journal.
7. CC Brachett, 2004. Journal of Pharmacotherapy.
8. M Gielen, 2002. Organometallic Chemistry.
9. DV Dick, W Rudolph, M Gielen, V. Wungerden, E. Kyra and N. Kees. 1998. Metal Based Drugs.
10. G.D Tiwari and M.N Mishra, 1980. Current Science.

- 
11. A Sigel and H Sigel, 2000. Indian J. Chem.
  12. H Singh, VK Srivastava, SN Shukla, and MK Srivastava, 2005. Indian J. Chem.
  13. HL Singh, S Varshney and AK Varshney, 1999. Appl. Organomet. Chem.
  14. Vinod Kumar and Rajesh Dhakarey, 2003. J. Ind. Council of Chemist.
  15. Alcock, JP Becker and AA Diamutis, 1992. J. Chem.
  16. William Herbert Gray, Gladwin Albert Hurst Buttle and Dora Stephenson, From the Welcome Chemical Research Laboratories, London (1937).
  17. S Baluja, A Solanki and N Kachhadia, 2006. J.Iranian Chem. Soc.
  18. NC Bharadwaj and RV Singh, 1994. Indian J. Chem.
  19. IM Kolthoff and R Elmquist, 1931. J. Am. Chem. Soc.
  20. K Chaturvedi, R Chaturvedi and N. Sharma, 2006. J. Ind. Chem. Soc.
  21. HL Singh, M Sharma and AK Varshney, Synth.1999. React. Met. Org. Chem.
  22. DM Adams, *Metal Ligands and Related Vibrations*, Arnold, London (1967).
  23. RM Silverstein, GC Bassler and TC Morrill, *Spectroscopic Identification of Organic Compounds*, John Willey and Sons, New York (1991).
  24. DE Ewing, Org. Magn. Reson., (1979).
  25. Shanu Paliwal, Gayatri Kumari and R Dhakarey, 2007. J. Indian Council Chem.